arXiv:1606.04622v1 [math.PR] 15 Jun 2016

On the last exit times for spectrally negative Lévy processes

Yingqiu Li[a], Chuancun Yin[b] and Xiaowen Zhou[c]

[a]School of Mathematics and Statistics, Changsha University of Science and Technology

[b]School of Statistics, Qufu Normal University

[c]Department of Mathematics and Statistics, Concordia University

Abstract

Using a new approach, for spectrally negative Lévy processes we find joint Laplace transforms involving the last exit time (from a semi-infinite interval), the value of the process at the last exit time and the associated occupation time, which generalizes some previous results.

Keywords: spectrally negative Lévy process, last exit time, Gerber-Shiu function, occupation time.
MSC classification: Primary 60G51; Secondary 60K30

1. Introduction

The first passage times, also known as the first exit times, have been studied extensively for stochastic processes. They find many applications in risk theory since the first exit time corresponds to the ruin time of a risk process. The joint distribution of quantities related to the ruin time, often referred as Gerber-Shiu functions, are very interested in the study of risk models. For spectrally negative Lévy processes those first passage time related quantities can often be expressed in terms of the associated scale functions.

The last passage time is also interested in risk theory. In a risk model, it can be regarded as the final recovery time after when there will be no more ruin; see [4] for early work in this respect. A Laplace transform for the last exit time from the positive part of the real line is first obtained in [3] for spectrally negative Lévy process using an identity between density of the first exit time and marginal density of the Lévy process. Applying the strong Markov property and the memoryless property for exponential random variables, in [2] Laplace transforms are also found for the last exit times (before an independent exponential time) from the positive part and the negative part of the real line, respectively. In both [3] and [2] the Laplace transforms are expressed in terms of the Laplace exponent and scale functions for the associated spectrally negative Lévy process.

In this paper we further consider the joint distribution concerning the last time when a spectrally negative Lévy process leaves a semi-infinite interval. More precisely, for a spectrally negative Lévy process running up to an exponential time, we find the joint Laplace transform of the occupation time until the last exit time, the last exit time and value of the process at the last exit time. They can be treated as Gerber-Shiu function like quantities for the last exit time of a spectrally negative Lévy process and generalize the previous results in [3] and [2]. Right before finalizing this paper we notice the recent work of [5] in which the joint distribution involving the the last exit time, the occupation time up to the last exit time and the value of the process prior to the last exit time. [5] adopts a time reversal argument and uses a previous result on occupation time.

Our approach is different from those in [3], [2] and [5]. We use the Poisson approach of [6] and [1] to find Laplace transforms of the occupation times. To overcome difficulties due to the non-stoping last exit time and the infinite activity of the Lévy processes, we adopt a perturbation approach, similar to that of [10], to approximate the last exit times by first exit times over certain events. We then apply the fluctuation theory for spectrally negative Lévy processes to express the approximating Laplace transforms in terms of scale functions and Laplace exponents before taking limits to reach the desired results. We also remark that whenever there is a jump at the last exit time, our result can not be directly obtained from that in [5] since the last exit time is not a stopping time although the approach of [5] can still be used to derive our results. We illustrate this with an example at the end of the paper. We also discuss the possible creeping behavior at the last exit time that is not covered in [5].

In the next section we present some preliminary results for spectrally negative Lévy processes. The main results and proofs are provided in Section 3. Some corollaries, discussions together with examples are also given in this section.

2. Preliminaries

In this section we collect some known fluctuation results for spectrally negative Lévy processes to prepare for the future proofs. Let process X be a spectrally negative Lévy process, i.e. a Lévy process with right continuous sample paths and with no positive jumps. Then for any $t > 0$,

$$\frac{1}{t}\log \mathbb{E}e^{\lambda X_t} = \psi(\lambda)$$

with

$$\psi(\lambda) = a\lambda + \frac{1}{2}\sigma^2\lambda^2 + \int_{-\infty}^{0}(e^{\lambda x} - 1 - \lambda x 1_{x>-1})\Pi(dx),$$

where Π is a σ-finite measure on $(-\infty, 0)$ satisfying $\int_{-\infty}^{0}(x^2 \wedge 1)\Pi(dx) < \infty$. Write $\Phi : [0, \infty) \mapsto [0, \infty)$ for the inverse function of ψ.

Scale functions play a key role in analyzing spectrally negative Lévy processes. The scale function $W^{(p)}$ for X is a nonnegative, increasing function specified by the following Laplace transform. For any $q \geq 0$,

$$\int_0^\infty e^{-\lambda x}W^{(q)}(x)dx = \frac{1}{\psi(\lambda) - q}, \;\; \lambda > \Phi(q),$$

and $W^{(q)}(x) := 0$ for $x < 0$. We also need the scale functions

$$Z^{(p)}(x) := 1 + p\int_0^x W^{(p)}(y)dy$$

and

$$Z^{(p)}(x, \theta) := e^{\theta x}\left(1 + (p - \psi(\theta))\int_0^x e^{-\theta y}W^{(p)}(y)dy\right).$$

with $Z^{(p)}(x) := 1$ and $Z^{(p)}(x, \theta) := e^{\theta x}$ for $x < 0$, respectively. Write $W(x), Z(x), Z(x, \theta)$ for $W^{(0)}(x), Z^{(0)}(x), Z^{(0)}(x, \theta)$, respectively.

For all $p, q \geq 0$, if X is of bounded variation, then $W^{(p)}(0) = W^{(q)}(0)$; if X is of unbounded variation, then

$$\lim_{\varepsilon \to 0+}\frac{W^{(p)}(\varepsilon)}{W^{(q)}(\varepsilon)} = 1;$$

see [8].

For any $a \in \mathbb{R}$, define one-sided exit times

$$\tau_a^+ := \inf\{t > 0 : X_t > a\}$$

and

$$\tau_a^- := \inf\{t > 0 : X_t < a\}$$

with the convention $\inf \emptyset = \infty$.

We collect some fluctuation identities for X concerning the solutions to the exit problems and potential measures, which can be found in [7] and [8]. For event A and random variable Y, write $\mathbb{E}[Y; A] = \mathbb{E}Y1_A$.

We first present solutions to the exit problems.

$$\mathbb{E}[e^{-q\tau_a^+}; \tau_a^+ < \infty] = e^{-\Phi(q)a},$$

$$\mathbb{E}_x[e^{-q\tau_0^-}; \tau_0^- < \infty] = Z^{(q)}(x) - \frac{q}{\Phi(q)} W^{(q)}(x),$$

$$\begin{aligned} \mathbb{E}_x[e^{-u\tau_0^- + vX_{\tau_0^-}}; \tau_0^- < \infty] &= Z^{(u)}(x, v) - \frac{u - \psi(v)}{\Phi(u) - v} W^{(u)}(x) \\ &= e^{vx}\left(1 + (u - \psi(v)) \int_0^x e^{-vy} W^{(u)}(y) \mathrm{d}y - \frac{u - \psi(v)}{\Phi(u) - v} e^{-vx} W^{(u)}(x)\right) \end{aligned} \tag{1}$$

and

$$\mathbb{E}_x[e^{-u\tau_0^- + \Phi(u)X_{\tau_0^-}}; \tau_0^- < \infty] = e^{\Phi(u)x} - \frac{1}{\Phi'(u)} W^{(u)}(x),$$

$$\begin{aligned} &\mathbb{E}_x[e^{-u\tau_0^- + vX_{\tau_0^-}}; \tau_0^- < \tau_a^+] \\ &= Z^{(u)}(x, v) - \frac{u - \psi(v)}{\Phi(u) - v} W^{(u)}(x) - \frac{W^{(u)}(x)}{W^{(u)}(a)} \left(Z^{(u)}(a, v) - \frac{u - \psi(v)}{\Phi(u) - v} W^{(u)}(a)\right) \\ &= Z^{(u)}(x, v) - \frac{W^{(u)}(x)}{W^{(u)}(a)} Z^{(u)}(a, v). \end{aligned} \tag{2}$$

We then present expressions for potential measures. For $a > 0$,

$$\int_0^\infty e^{-qt} \mathbb{P}\{X_t \in dy, t < \tau_a^+\} dt = \left(e^{-\Phi(q)a} W^{(q)}(a - y) - W^{(q)}(-y)\right) dy. \tag{3}$$

For $a \leq 0$,

$$\int_0^\infty e^{-qt} \mathbb{P}\{X_t \in dy, t < \tau_a^-\} dt = \left(e^{-\Phi(q)(y-a)} W^{(q)}(-a) - W^{(q)}(-y)\right) dy. \tag{4}$$

For an independent exponential random variable e_r with rate r, let

$$T^+(r) := \sup\{0 < t \leq e_r : X_t < 0\}$$

and

$$T^-(r) := \sup\{0 < t \leq e_r : X_t > 0\}$$

with the convention $\sup \emptyset = 0$. Then $T^+(r)$ and $T^-(r)$ are, respectively, the last times of exiting the negative and positive part of the real line up to the time e_r. Write T^+ and T^- for $T^+(0)$ and $T^-(0)$, respectively.

In this paper, for constants $p, q, \theta, r > 0$ and $x \in \mathbb{R}$, we want to find expressions for

$$\mathbb{E}_x[e^{-pT^+(r) - q\int_0^{T^+(r)} 1_{X_s < 0} ds + \theta X(T^+(r))}; T^+(r) = e_r],$$

$$\mathbb{E}_x[e^{-pT^+(r)-q\int_0^{T^+(r)}1_{X_s<0}ds};T^+(r)<e_r]$$

and

$$\mathbb{E}_x[e^{-pT^-(r)+\theta X(T^-(r))-q\int_0^{T^-(r)}1_{X_s>0}ds};T^-(r)<e_r],$$
$$\mathbb{E}_x[e^{-pT^-(r)-q\int_0^{T^-(r)}1_{X_s>0}ds-\theta X(T^-(r))};T^-(r)=e_r].$$

3. Main Results

We first proceed to find expressions for

$$\omega_1^+(x):=\mathbb{E}_x[e^{-pT^+(r)-q\int_0^{T^+(r)}1_{X_s\le 0}ds};T^+(r)<e_r]$$

and

$$\omega_2^+(x):=\mathbb{E}_x[e^{-pT^+(r)-q\int_0^{T^+(r)}1_{X_s\le 0}ds+\theta X(T^+(r))};T^+(r)=e_r].$$

Theorem 3.1. *For any $p,q,r,\theta>0$ and $x\in\mathbb{R}$, we have*

$$\begin{aligned}\omega_1^+(x)=&\frac{r(W^{(r)}(x)-W^{(p+r)}(x))}{\Phi(r)}+\frac{r(\Phi(p+q+r)-\Phi(p+r))}{q\Phi(r)}Z^{(p+r)}(x,\Phi(p+q+r))\\&-r\int_0^x W^{(r)}(y)dy\end{aligned}\tag{5}$$

and

$$\begin{aligned}\omega_2^+(x)=&\frac{r}{p+q+r-\psi(\theta)}\left(-\frac{p+r-\psi(\theta)}{q}\times\frac{\Phi(p+q+r)-\Phi(p+r)}{\Phi(p+r)}\right)Z^{(p+r)}(x,\Phi(p+q+r))\\&+\frac{r}{p+q+r-\psi(\theta)}Z^{(p+r)}(x,\theta).\end{aligned}\tag{6}$$

Proof. The proof is carried out in two steps. We first assume that X is of bounded variation. Write e_q for an independent exponential random variable. Let $T_1<T_2<\dots$ be the sequence of arrival times for an independent Poisson process with rate q. Let

$$A^+:=\cap_{i:T_i<T^+(r)}\{X_{T_i}>0\}.$$

Then

$$\omega_1^+(x)=\mathbb{E}_x\left[e^{-pT^+(r)};T^+(r)<e_r,A^+\right]$$

and

$$\omega_2^+(x)=\mathbb{E}_x\left[e^{-pT^+(r)+\theta X(e_r)};T^+(r)=e_r,A^+\right]$$

by a property of Poisson process; see a similar argument in [6]. Then conditioning on T_1, by the the memoryless property for exponential distribution and the strong Markov property,

$$\begin{aligned}\omega_1^+(0)&=\mathbb{P}\{\tau_0^-<e_r\}+\int_{-\infty}^0\mathbb{E}[e^{-p\tau_0^-};\tau_0^-<e_r,X_{\tau_0^-}\in dy]\mathbb{E}_y[e^{-p\tau_0^+};\tau_0^+<e_q\wedge e_r]\omega_1^+(0)\\&=1-\mathbb{E}e^{-r\tau_0^-}+\int_{-\infty}^0\mathbb{E}[e^{-(p+r)\tau_0^-};X_{\tau_0^-}\in dy]\mathbb{E}_y e^{-(p+q+r)\tau_0^+}\omega_1^+(0)\\&=\frac{r}{\Phi(r)}W^{(r)}(0)+\mathbb{E}e^{-(p+r)\tau_0^-+\Phi(p+q+r)X(\tau_0^-)}\omega_1^+(0)\\&=\frac{r}{\Phi(r)}W^{(r)}(0)+\left(1-\frac{q}{\Phi(p+q+r)-\Phi(p+r)}W^{(p+r)}(0)\right)\omega_1^+(0),\end{aligned}\tag{7}$$

where we have used the identity (1). Solving (7) for $\omega_1^+(0)$ we have

$$\omega_1^+(0) = \frac{r}{q\Phi(r)}(\Phi(p+q+r) - \Phi(p+r)). \tag{8}$$

Similarly,

$$\begin{aligned}
\omega_2^+(0) &= \int_{-\infty}^0 \mathbb{E}[e^{-p\tau_0^-}; \tau_0^- < e_r, X_{\tau_0^-} \in dy] \\
&\qquad \times \left(\mathbb{E}_y[e^{-p\tau_0^+}; \tau_0^+ < e_q \wedge e_r]\omega_2^+(0) + \mathbb{E}_y[e^{-pe_r+\theta X(e_r)}; e_r < \tau_0^+ \wedge e_q]\right) \\
&= \mathbb{E}e^{-(p+r)\tau_0^- + \Phi(p+q+r)X(\tau_0^-)}\omega_2^+(0) \\
&\quad + \frac{r}{p+q+r-\psi(\theta)}\left(\mathbb{E}e^{-(p+r)\tau_0^- + \theta X(\tau_0^-)} - \mathbb{E}e^{-(p+r)\tau_0^- + \Phi(p+q+r)X(\tau_0^-)}\right) \\
&= \left(1 - \frac{q}{\Phi(p+q+r) - \Phi(p+r)}W^{(p+r)}(0)\right)\omega_2^+(0) \\
&\quad + \frac{r}{p+q+r-\psi(\theta)}\left(\frac{q}{\Phi(p+q+r) - \Phi(p+r)}W^{(p+r)}(0) - \frac{p+r-\psi(\theta)}{\Phi(p+r)-\theta}W^{(p+r)}(0)\right),
\end{aligned} \tag{9}$$

where for $y < 0$ and $\theta > p+q+r$, by identity (3) we have

$$\begin{aligned}
\mathbb{E}_y[e^{-pe_r+\theta X(e_r)}; e_r < \tau_0^+ \wedge e_q] &= \mathbb{E}_y[e^{-(p+q)e_r+\theta X(e_r)}; e_r < \tau_0^+] \\
&= \int_{-\infty}^0 e^{\theta z} r\left(e^{\Phi(p+q+r)y}W^{(p+q+r)}(-z) - W^{(p+q+r)}(y-z)\right) \\
&= e^{\Phi(p+q+r)y}\frac{r}{\psi(\theta) - p - q - r} - e^{\theta y}\frac{r}{\psi(\theta) - p - q - r}.
\end{aligned}$$

Solving (9) for $\omega_2^+(0)$ we have

$$\omega_2^+(0) = \frac{r}{p+q+r-\psi(\theta)}\left(1 - \frac{p+r-\psi(\theta)}{q} \times \frac{\Phi(p+q+r) - \Phi(p+r)}{\Phi(p+r)-\theta}\right). \tag{10}$$

We now consider process X with sample paths of unbounded variation. For any $\epsilon > 0$, let

$$T_\varepsilon^+(r) := \inf\{T^+(r) \le s \le e_r : X_s > \varepsilon\}$$

with the convention $\inf\emptyset = e_r$. Write $T_\varepsilon^+ = T_\varepsilon^+(0)$. Then either $T^+(r) < e_r$ or $T^+(r) = e_r$, we have $T_\varepsilon^+(r) \downarrow T^+(r)$ as $\varepsilon \to 0+$. Write

$$\omega_{1,\varepsilon}^+(x) := \mathbb{E}_x[e^{-pT_\varepsilon^+(r) - q\int_0^{T_\varepsilon^+(r)} 1_{X_s \le 0} ds}; T_\varepsilon^+(r) < e_r]$$

and

$$\omega_{2,\varepsilon}^+(x) := \mathbb{E}_x[e^{-pT_\varepsilon^+(r) - q\int_0^{T_\varepsilon^+(r)} 1_{X_s \le 0} ds + \theta X(e_r)}; T_\varepsilon^+(r) = e_r].$$

Then for

$$\begin{aligned}
A_\varepsilon^+ &:= \cap_{i:T_i < T_\varepsilon^+(r)}\{X_{T_i} > 0\}, \\
\omega_{1,\varepsilon}^+(x) &= \mathbb{E}_x\left[e^{-pT_\varepsilon^+(r)}; T_\varepsilon^+(r) < e_r, A_\varepsilon^+\right], \\
\omega_{2,\varepsilon}^+(x) &= \mathbb{E}_x\left[e^{-pT_\varepsilon^+(r) + \theta X(e_r)}; T_\varepsilon^+(r) = e_r, A_\varepsilon^+\right],
\end{aligned}$$

by the property of Poisson process. It follows that

$$\lim_{\varepsilon\to 0+}\omega^+_{i,\varepsilon}(x)=\omega^+_i(x),\quad i=1,2.$$

Notice that by (3)

$$\begin{aligned}&\mathbb{E}[e^{-pT^+_\varepsilon(r)};T^+_\varepsilon(r)<e_r,T_1<\tau^+_\varepsilon\wedge e_r,A^+]\\&\le\mathbb{P}\{T_1<\tau^+_\varepsilon,0\le X_{T_1}\le\varepsilon\}\\&=o(W(\varepsilon)).\end{aligned}\tag{11}$$

Comparing the first Poisson arrival time T_1 with τ^+_ε and e_r we have

$$\begin{aligned}\omega^+_{1,\varepsilon}(0)&=\mathbb{E}[e^{-p\tau^+_\varepsilon};\tau^+_\varepsilon<e_q\wedge e_r]\left(\mathbb{P}_\varepsilon[\tau^-_0>e_r\}\right.\\&\qquad\left.+\int_{-\infty}^0\mathbb{E}_\varepsilon[e^{-p\tau^-_0};\tau^-_0<e_r,X_{\tau^-_0}\in dy]\mathbb{E}_y[e^{-p\tau^+_0};\tau^+_0<e_q\wedge e_r]\omega^+_{1,\varepsilon}(0)\right)\\&\quad+o(W(\varepsilon))\\&=e^{-\Phi(p+q+r)\varepsilon}\left\{1-Z^{(r)}(\varepsilon)+\frac{r}{\Phi(r)}W^{(r)}(\varepsilon)+\mathbb{E}_\varepsilon e^{-(p+r)\tau^-_0+\Phi(p+q+r)X(\tau^-_0)}\omega^+_{1,\varepsilon}(0)\right\}\\&\quad+o(W(\varepsilon))\\&=e^{-\Phi(p+q+r)\varepsilon}\frac{r}{\Phi(r)}W^{(r)}(\varepsilon)+o(W(\varepsilon))\\&\quad+\left(1-\frac{q}{\Phi(p+q+r)-\Phi(p+r)}e^{-\Phi(p+q+r)\varepsilon}W^{(p+r)}(\varepsilon)\right)\omega^+_{1,\varepsilon}(0).\end{aligned}\tag{12}$$

Solve (12) for $\omega^+_\varepsilon(0)$ and take a limit. Then

$$\begin{aligned}\omega^+_1(0)&=\lim_{\varepsilon\to 0+}\omega^+_{1,\varepsilon}(0)\\&=\lim_{\varepsilon\to 0+}\frac{e^{-\Phi(p+q+r)\varepsilon}\frac{r}{\Phi(r)}W^{(r)}(\varepsilon)}{\frac{q}{\Phi(p+q+r)-\Phi(p+r)}e^{-\Phi(p+q+r)\varepsilon}W^{(p+r)}(\varepsilon)}\\&=\frac{r}{q\Phi(r)}(\Phi(p+q+r)-\Phi(p+r)).\end{aligned}$$

Similarly, for $\theta>0$,

$$\begin{aligned}&\mathbb{E}[e^{-pT^+_\varepsilon(r)+\theta X(T^+_\varepsilon(r))};T^+_\varepsilon(r)=e_r,T_1<\tau^+_\varepsilon\wedge e_r,A^+_\varepsilon]\\&\le\mathbb{P}\{T_1<\tau^+_\varepsilon,0\le X_{T_1}\le\varepsilon\}\\&=o(W(\varepsilon)).\end{aligned}$$

Then for $\theta > p+q+r$,

$$
\begin{aligned}
\omega^+_{2,\varepsilon}(0) &= \mathbb{E}[e^{-p\tau^+_\varepsilon}; \tau^+_\varepsilon < e_q \wedge e_r] \int_{-\infty}^0 \mathbb{E}_\varepsilon[e^{-p\tau^-_0}; \tau^-_0 < e_r, X_{\tau^-_0} \in dy] \\
&\quad \times \left(\mathbb{E}_y[e^{-p\tau^+_0}; \tau^+_0 < e_q \wedge e_r] \omega^+_{2,\varepsilon}(0) + \mathbb{E}_y[e^{-pe_r+\theta X(e_r)}; e_r < \tau^+_0 \wedge e_q] \right) \\
&\quad + \mathbb{E}[e^{-pe_r+\theta X(e_r)}; e_r < \tau^+_\varepsilon \wedge e_q, X_{e_r} < 0] + o(W(\varepsilon)) \\
&= e^{-\Phi(p+q+r)\varepsilon} \mathbb{E}_\varepsilon e^{-(p+r)\tau^-_0 + \Phi(p+q+r)X(\tau^-_0)} \omega^+_{2,\varepsilon}(0) \\
&\quad + e^{-\Phi(p+q+r)\varepsilon} \frac{r}{p+q+r-\psi(\theta)} \left(\mathbb{E}_\varepsilon e^{-(p+r)\tau^-_0 + \theta X(\tau^-_0)} - \mathbb{E}_\varepsilon e^{-(p+r)\tau^-_0 + \Phi(p+q+r)X(\tau^-_0)} \right) \\
&\quad + \frac{r}{p+q+r-\psi(\theta)} (1 - e^{(\theta - \Phi(p+q+r))\varepsilon}) + o(W(\varepsilon)) \\
&= \left(1 - \frac{q}{\Phi(p+q+r) - \Phi(p+r)} e^{-\Phi(p+q+r)\varepsilon} W^{(p+r)}(\varepsilon) \right) \omega^+_{2,\varepsilon}(0) + o(W(\varepsilon)) \\
&\quad + \frac{r}{p+q+r-\psi(\theta)} \left(\frac{q}{\Phi(p+q+r) - \Phi(p+r)} W^{(p+r)}(\varepsilon) \right. \\
&\qquad \left. - e^{(\theta - \Phi(p+q+r))\varepsilon} \frac{p+r-\psi(\theta)}{\Phi(p+r) - \theta} W^{(p+r)}(\varepsilon) \right),
\end{aligned} \tag{13}
$$

where by (3)

$$
\begin{aligned}
&\mathbb{E}[e^{-pe_r+\theta X(e_r)}; e_r < \tau^+_\varepsilon \wedge e_q, X_{e_r} < 0] \\
&= \mathbb{E}[e^{-(p+q)e_r+\theta X(e_r)}; e_r < \tau^+_\varepsilon] \\
&= \int_{-\infty}^0 e^{\theta y} r \left(e^{-\Phi(p+q+r)\varepsilon} W^{(p+q+r)}(\varepsilon - y) - W^{(p+q+r)}(-y) \right) dy \\
&= r e^{-\Phi(p+q+r)\varepsilon} \int_{-\infty}^{-\varepsilon} e^{\theta(y+\varepsilon)} W^{(p+q+r)}(-y) dy - r \int_{-\infty}^0 e^{\theta y} W^{(p+q+r)}(-y) dy \\
&= e^{(\theta - \Phi(p+q+r))\varepsilon} \frac{r}{\psi(\theta) - p - q - r} - \frac{r}{\psi(\theta) - p - q - r} + o(W(\varepsilon)).
\end{aligned}
$$

Then

$$
\begin{aligned}
\omega^+_2(0) &= \lim_{\varepsilon \to 0+} \omega^+_{2,\varepsilon}(0) \\
&= \frac{\frac{r}{p+q+r-\psi(\theta)} \left(\frac{q}{\Phi(p+q+r) - \Phi(p+r)} - e^{(\theta - \Phi(p+q+r))\varepsilon} \frac{p+r-\psi(\theta)}{\Phi(p+r) - \theta} \right)}{\frac{q}{\Phi(p+q+r) - \Phi(p+r)}} \\
&= \frac{r}{p+q+r-\psi(\theta)} \left(1 - \frac{p+r-\psi(\theta)}{q} \times \frac{\Phi(p+q+r) - \Phi(p+r)}{\Phi(p+r) - \theta} \right).
\end{aligned} \tag{14}
$$

For $x < 0$,

$$
\begin{aligned}
\omega^+_1(x) &= \mathbb{E}_x[e^{-(p+q)\tau^+_0}; \tau^+_0 < e_r] \omega^+_1(0) \\
&= e^{\Phi(p+q+r)x} \frac{r}{q\Phi(r)} (\Phi(p+q+r) - \Phi(p+r)).
\end{aligned}
$$

Similarly,

$$\begin{aligned}\omega_2^+(x) &= \mathbb{E}_x[e^{-(p+q)e_r+\theta X(e_r)}; e_r < \tau_0^+] + \mathbb{E}_x[e^{-(p+q)\tau_0^+}; \tau_0^+ < e_r]\omega_2^+(0) \\ &= \frac{r}{p+q+r-\psi(\theta)}\left(e^{\theta x} - e^{\Phi(p+q+r)x}\frac{p+r-\psi(\theta)}{q} \times \frac{\Phi(p+q+r)-\Phi(p+r)}{\Phi(p+r)-\theta}\right).\end{aligned} \tag{15}$$

By analytical extension, (15) holds for all $\theta \geq 0$.

For $x > 0$,

$$\begin{aligned}\omega_1^+(x) &= \mathbb{P}_x\{\tau_0^- > e_r\} + \int_{-\infty}^0 \mathbb{E}_x[e^{-p\tau_0^-}; \tau_0^- < e_r, X_{\tau_0^-} \in dy]\mathbb{E}_y[e^{-(p+q)\tau_0^+}; \tau_0^+ < e_r]\omega_1^+(0) \\ &= \frac{r}{\Phi(r)}W^{(r)}(x) - r\int_0^x W^{(r)}(y)dy \\ &\quad + \frac{r(\Phi(p+q+r)-\Phi(p+r))}{q\Phi(r)}\left\{Z^{(p+r)}(x,\Phi(p+q+r)) - \frac{q}{\Phi(p+q+r)-\Phi(p+r)}W^{(p+r)}(x)\right\} \\ &= \frac{r(W^{(r)}(x)-W^{(p+r)}(x))}{\Phi(r)} - r\int_0^x W^{(r)}(y)dy \\ &\quad + \frac{r(\Phi(p+q+r)-\Phi(p+r))}{q\Phi(r)}Z^{(p+r)}(x,\Phi(p+q+r))\end{aligned}$$

and

$$\begin{aligned}\omega_2^+(x) &= \int_{-\infty}^0 \mathbb{E}_x[e^{-p\tau_0^-}; \tau_0^- < e_r, X_{\tau_0^-} \in dy] \\ &\qquad \times \left(\mathbb{E}_y[e^{-(p+q)\tau_0^+}; \tau_0^+ < e_r]\omega_2^+(0) + \mathbb{E}_y[e^{-(p+q)e_r+\theta X(e_r)}; e_r < \tau_0^+]\right) \\ &= \mathbb{E}_x e^{-(p+r)\tau_0^- + \Phi(p+q+r)X(\tau_0^-)}\omega_2^+(0) \\ &\quad + \frac{r}{p+q+r-\psi(\theta)}\left(\mathbb{E}_x e^{-(p+r)\tau_0^- + \theta X(\tau_0^-)} - \mathbb{E}_x e^{-(p+r)\tau_0^- + \Phi(p+q+r)X(\tau_0^-)}\right) \\ &= \frac{r}{p+q+r-\psi(\theta)}\left(-\frac{p+r-\psi(\theta)}{q} \times \frac{\Phi(p+q+r)-\Phi(p+r)}{\Phi(p+r)-\theta}\right)Z^{(p+r)}(x,\Phi(p+q+r)) \\ &\quad + \frac{r}{p+q+r-\psi(\theta)}Z^{(p+r)}(x,\theta).\end{aligned}$$

The desired results then follow. □

Remark 3.1. *For $x < 0$,*

$$\begin{aligned}\mathbb{E}_x e^{-pT^+(r)} &= \lim_{q,\theta\to 0+}(\omega_1^+(0) + \omega_2^+(0)) \\ &= \frac{r}{p+r} + re^{\Phi(p+r)x}\Phi'(p+r)\left(\frac{1}{\Phi(r)} - \frac{1}{\Phi(p+r)}\right).\end{aligned} \tag{16}$$

For $x > 0$,

$$\begin{aligned}\mathbb{E}_x e^{-pT^+(r)} &= \frac{r(W^{(r)}(x)-W^{(p+r)}(x))}{\Phi(r)} + re^{\Phi(p+r)x}\Phi'(p+r)\left(\frac{1}{\Phi(r)} - \frac{1}{\Phi(p+r)}\right) \\ &\quad + 1 - Z^{(r)}(x) + \frac{r}{p+r}Z^{(p+r)}(x),\end{aligned} \tag{17}$$

which recovers Theorem 2 in [2].

Remark 3.2. *For $\psi'(0+) > 0$, letting $r, q \to 0+$ we have*

$$\mathbb{E}_x e^{-pT^+} = \psi'(0+)(W(x) - W^{(p)}(x)) + e^{\Phi(p)x}\psi'(0+)\Phi'(p),$$

which recovers a result in [3]*; letting $r, p \to 0+$ for $\psi'(0+) > 0$, we have*

$$\mathbb{E}_x e^{-q\int_0^\infty 1_{X_s<0}ds} = \frac{\psi'(0+)}{q}\Phi(q)Z(x, \Phi(q)),$$

which recovers Corollary 3 (ii) of [9].

We also want to find

$$\omega_1^-(x) := \mathbb{E}_x[e^{-pT^-(r)+\theta X_{T^-(r)}-q\int_0^{T^-(r)} 1_{X_s\geq 0}ds}; T^-(r) < e_r]$$

and

$$\omega_2^-(x) := \mathbb{E}_x[e^{-pT^-(r)-q\int_0^{T^-(r)} 1_{X_s\geq 0}ds-\theta X(T^-(r))}; T^-(r) = e_r].$$

Theorem 3.2. *For any $p, q, r, \theta > 0$ and $x \in \mathbb{R}$, we have*

$$\begin{aligned}\omega_1^-(x) = {} & \frac{1}{q}(\Phi(p+q+r) - \Phi(p+r))Z^{(p+q+r)}(x, \Phi(p+r)) \\ & \times \left(\frac{p+q+r-\psi(\theta+\Phi(r))}{\Phi(p+q+r)-\theta-\Phi(r)} - \frac{p+q+r-\psi(\theta)}{\Phi(p+q+r)-\theta}\right) \\ & + Z^{(p+q+r)}(x, \theta) - Z^{(p+q+r)}(x, \theta+\Phi(r))\end{aligned} \tag{18}$$

and

$$\omega_2^-(x) = \frac{r}{q} \times \frac{\Phi(p+q+r)-\Phi(p+r)}{\theta+\Phi(p+q+r)}Z^{(p+q+r)}(x, \Phi(p+r)) - r\int_0^x e^{-\theta y}W^{(p+q+r)}(x-y)dy. \tag{19}$$

Proof. If X is of bounded variation, for

$$A^- := \cap_{i:T_i<T^-(r)}\{X_{T_i} < 0\},$$

we have

$$\omega_1^-(x) = \mathbb{E}_x\left[e^{-pT^-(r)+\theta X_{T^-(r)}-q\int_0^{T^-(r)} 1_{X_s\geq 0}ds}; T^-(r) < e_r, A^-\right].$$

Then

$$\begin{aligned}\omega_1^-(0) = {} & \int_{-\infty}^0 \mathbb{E}[e^{-p\tau_0^-}; \tau_0^- < e_q \wedge e_r, X_{\tau_0^-} \in dy] \\ & \times \left(\mathbb{E}_y[e^{-p\tau_0^+}; \tau_0^+ < e_r]\omega_1^-(0) + e^{\theta y}\mathbb{P}_y\{e_r < \tau_0^+\}\right) \\ = {} & \int_{-\infty}^0 \mathbb{E}[e^{-(p+q+r)\tau_0^-}; X_{\tau_0^-} \in dy]\left(\mathbb{E}_y e^{\Phi(p+r)y}\omega_1^-(0) + e^{\theta y} - e^{(\theta+\Phi(r))y}\right) \qquad (20) \\ = {} & \left(1 - \frac{qW^{(p+q+r)}(0)}{\Phi(p+q+r)-\Phi(p+r)}\right)\omega_1^-(0) - \frac{p+q+r-\psi(\theta)}{\Phi(p+q+r)-\theta}W^{(p+q+r)}(0) \\ & + \frac{p+q+r-\psi(\theta+\Phi(r))}{\Phi(p+q+r)-\theta-\Phi(r)}W^{(p+q+r)}(0).\end{aligned}$$

Then

$$\omega_1^-(0) = \frac{1}{q}(\Phi(p+q+r) - \Phi(p+r)) \left(\frac{p+q+r-\psi(\theta+\Phi(r))}{\Phi(p+q+r)-\theta-\Phi(r)} - \frac{p+q+r-\psi(\theta)}{\Phi(p+q+r)-\theta} \right).$$

Similarly,

$$\begin{aligned}
\omega_2^-(0) &= \mathbb{E}[e^{-pe_r-\theta X(e_r)}; e_r < \tau_0^- \wedge e_q] \\
&\quad + \int_{-\infty}^0 \mathbb{E}[e^{-p\tau_0^-}; \tau_0^- < e_q \wedge e_r, X_{\tau_0^-} \in dy] \mathbb{E}_y[e^{-p\tau_0^+}; \tau_0^+ < e_r] \omega_1^-(0) \\
&= \mathbb{E}[e^{-(p+q)e_r-\theta X(e_r)}; e_r < \tau_0^-] + \mathbb{E}e^{-(p+q+r)\tau_0^- + \Phi(p+r)\tau_0^-} \\
&= \frac{r}{\theta+\Phi(p+q+r)} W^{(p+q+r)}(0) + \left(1 - \frac{qW^{(p+q+r)}(0)}{\Phi(p+q+r)-\Phi(p+r)}\right) \omega_2^-(0).
\end{aligned}$$

where by identity (4)

$$\begin{aligned}
&\mathbb{E}[e^{-(p+q)e_r-\theta X(e_r)}; e_r < \tau_0^-] \\
&= r \int_0^\infty e^{-\theta y} \int_0^\infty \mathbb{E}[e^{-(p+q+r)t}; t < \tau_0^-, X_t \in dy] \\
&= r \int_0^\infty e^{-\theta y - \Phi(p+q+r)y} W^{(p+q+r)}(0) dy \\
&= \frac{r}{\theta+\Phi(p+q+r)} W^{(p+q+r)}(0).
\end{aligned}$$

Then

$$\omega_2^-(0) = \frac{r}{q} \times \frac{\Phi(p+q+r)-\Phi(p+r)}{\theta+\Phi(p+q+r)}.$$

To deal with process X with sample paths of unbounded variation, for $\varepsilon > 0$ and $x \in \mathbb{R}$, let

$$T_\varepsilon^-(r) := \inf\{T^-(r) \le s \le e_r : X_s < -\varepsilon\}$$

with the convention $\inf \emptyset = e_r$ and

$$\omega_{1,\varepsilon}^-(x) := \mathbb{E}_x[e^{-pT_\varepsilon^-(r) + \theta X_{T_\varepsilon^-(r)} - q\int_0^{T_\varepsilon^-(r)} 1_{X_s \ge 0} ds}; T_\varepsilon^-(r) < e_q],$$

$$\omega_{2,\varepsilon}^-(x) := \mathbb{E}_x[e^{-pT_\varepsilon^-(r) - q\int_0^{T_\varepsilon^-(r)} 1_{X_s \ge 0} ds - \theta X(e_r)}; T_\varepsilon^-(r) = e_q].$$

Then for

$$A_\varepsilon^- := \cap_{i: T_i < T_\varepsilon^-(r)} \{X_{T_i} < 0\},$$

we have

$$\omega_{1,\varepsilon}^-(x) = \mathbb{E}_x \left[e^{-pT_\varepsilon^-(r) + \theta X_{T_\varepsilon^-(r)} - q\int_0^{T_\varepsilon^-(r)} 1_{X_s \ge 0} ds}; T_\varepsilon^-(r) < e_r, A_\varepsilon^- \right].$$

By (4),

$$\begin{aligned}
&\mathbb{E}[e^{-pT_\varepsilon^-(r) + \theta X(T_\varepsilon^-(r))}; T_1 < e_r \wedge \tau_{-\varepsilon}^-, T_\varepsilon^-(r) < e_r, A_\varepsilon^-] \\
&\le \mathbb{P}\{T_1 < \tau_{-\varepsilon}^-, -\varepsilon \le X_{T_1} \le 0\} \\
&= o(W(\varepsilon)).
\end{aligned}$$

Then

$$\begin{aligned}
\omega_{1,\varepsilon}^-(0) &= \int_{-\infty}^{-\varepsilon} \mathbb{E}[e^{-p\tau_{-\varepsilon}^- + \theta y}; \tau_{-\varepsilon}^- < e_q \wedge e_r, X_{\tau_{-\varepsilon}^-} \in dy] \mathbb{P}_y\{\tau_0^+ > e_r\} \\
&\quad + \int_{-\infty}^{-\varepsilon} \mathbb{E}[e^{-p\tau_{-\varepsilon}^-}; \tau_{-\varepsilon}^- < e_q \wedge e_r, X_{\tau_{-\varepsilon}^-} \in dy] \mathbb{E}_y[e^{-p\tau_0^+}; \tau_0^+ < e_r] \omega_{1,\varepsilon}^-(0) \\
&\quad + o(W(\varepsilon)) \\
&= \int_{-\infty}^{-\varepsilon} \mathbb{E}[e^{-(p+q+r)\tau_{-\varepsilon}^- + \theta y}; X_{\tau_{-\varepsilon}^-} \in dy](1 - e^{\Phi(r)y}) \\
&\quad + \int_{-\infty}^{-\varepsilon} \mathbb{E}[e^{-(p+q+r)\tau_{-\varepsilon}^-}; X_{\tau_{-\varepsilon}^-} \in dy] e^{\Phi(p+r)y} \omega_{1,\varepsilon}^-(0) + o(W(\varepsilon)) \\
&= 1 + o(W(\varepsilon)) - \frac{p+q+r-\psi(\theta)}{\Phi(p+q+r)-\theta} e^{-\theta\varepsilon} W^{(p+q+r)}(\varepsilon) \\
&\quad - 1 + o(W(\varepsilon)) + \frac{p+q+r-\psi(\theta+\Phi(r))}{\Phi(p+q+r)-\theta-\Phi(r)} e^{-(\theta+\Phi(r))\varepsilon} W^{(p+q+r)}(\varepsilon) \\
&\quad + \left(1 - \frac{q}{\Phi(p+q+r)-\Phi(p+r)} e^{-\Phi(p+r)\varepsilon} W^{(p+q+r)}(\varepsilon)\right) \omega_{1,\varepsilon}^-(0).
\end{aligned}$$

Then

$$\begin{aligned}
\omega_1^-(0) &= \lim_{\varepsilon \to 0+} \omega_{1,\varepsilon}^-(0) \\
&= \frac{1}{q}(\Phi(p+q+r) - \Phi(p+r)) \left(\frac{p+q+r-\psi(\theta+\Phi(r))}{\Phi(p+q+r)-\theta-\Phi(r)} - \frac{p+q+r-\psi(\theta)}{\Phi(p+q+r)-\theta} \right).
\end{aligned}$$

Similarly,

$$\begin{aligned}
\omega_{2,\varepsilon}^-(0) &= \int_{-\infty}^{-\varepsilon} \mathbb{E}[e^{-p\tau_{-\varepsilon}^-}; \tau_{-\varepsilon}^- < e_q \wedge e_r, X_{\tau_{-\varepsilon}^-} \in dy] \mathbb{E}_y[e^{-p\tau_0^+}; \tau_0^+ < e_r] \omega_{2,\varepsilon}^-(0) \\
&\quad + \mathbb{E}[e^{-pe_r - \theta X(e_r)}; e_r < \tau_{-\varepsilon}^- \wedge e_q] + o(W(\varepsilon)) \\
&= \left(1 - \frac{q}{\Phi(p+q+r)-\Phi(p+r)} e^{-\Phi(p+r)\varepsilon} W^{(p+q+r)}(\varepsilon)\right) \omega_{2,\varepsilon}^-(0) \\
&\quad + \frac{r}{\theta+\Phi(p+q+r)} W^{(p+q+r)}(\varepsilon) + o(W(\varepsilon)),
\end{aligned}$$

where

$$\begin{aligned}
&\mathbb{E}[e^{-pe_r - \theta X(e_r)}; e_r < \tau_{-\varepsilon}^- \wedge e_q] \\
&= r \int_{-\varepsilon}^{\infty} e^{-\theta y} \left(e^{-\Phi(p+q+r)(y+\varepsilon)} W^{(p+q+r)}(\varepsilon) - W^{(p+q+r)}(-y) \right) dy \\
&= \frac{r}{\theta+\Phi(p+q+r)} W^{(p+q+r)}(\varepsilon) + o(W(\varepsilon)).
\end{aligned}$$

Then

$$\omega_2^-(0) = \frac{r}{q} \times \frac{\Phi(p+q+r)-\Phi(p+r)}{\theta+\Phi(p+q+r)}.$$

For $x<0$,

$$\begin{aligned}\omega_1^-(x) &= e^{\theta x}\mathbb{P}_x\{e_r<\tau_0^+\}+\mathbb{E}_x[e^{-p\tau_0^+};\tau_0^+<e_r]\omega_1^-(0)\\ &= e^{\theta x}(1-e^{\Phi(r)x})\\ &\quad+\frac{e^{\Phi(p+r)x}}{q}(\Phi(p+q+r)-\Phi(p+r))\left(\frac{p+q+r-\psi(\theta+\Phi(r))}{\Phi(p+q+r)-\theta-\Phi(r)}-\frac{p+q+r-\psi(\theta)}{\Phi(p+q+r)-\theta}\right).\end{aligned}$$

$$\begin{aligned}\omega_2^-(x) &= \mathbb{E}_x[e^{-p\tau_0^+};\tau_0^+<e_r]\omega_2^-(0)\\ &= \frac{re^{\Phi(p+r)x}}{q}\times\frac{\Phi(p+q+r)-\Phi(p+r)}{\theta+\Phi(p+q+r)}.\end{aligned}$$

For $x>0$,

$$\begin{aligned}\omega_1^-(x) &= \int_{-\infty}^0\mathbb{E}_x[e^{-(p+q)\tau_0^-};\tau_0^-<e_r,X_{\tau_0^-}\in dy]\left(\mathbb{E}_y[e^{-p\tau_0^+};\tau_0^+<e_r]\omega_1^-(0)+e^{\theta y}\mathbb{P}_y\{e_r<\tau_0^+\}\right)\\ &= \mathbb{E}_xe^{-(p+q+r)\tau_0^-+\Phi(p+r)X(\tau_0^-)}\omega_1^-(0)+\mathbb{E}_xe^{-(p+q+r)\tau_0^-+\theta X(\tau_0^-)}-\mathbb{E}_xe^{-(p+q+r)\tau_0^-+(\theta+\Phi(r))X(\tau_0^-)}\\ &= \left\{Z^{(p+q+r)}(x,\Phi(p+r))-\frac{q}{\Phi(p+q+r)-\Phi(p+r)}W^{(p+q+r)}(x)\right\}\\ &\quad\times\frac{1}{q}(\Phi(p+q+r)-\Phi(p+r))\left(\frac{p+q+r-\psi(\theta+\Phi(r))}{\Phi(p+q+r)-\theta-\Phi(r)}-\frac{p+q+r-\psi(\theta)}{\Phi(p+q+r)-\theta}\right)\\ &\quad+Z^{(p+q+r)}(x,\theta)-\frac{p+q+r-\psi(\theta)}{\Phi(p+q+r)-\theta}W^{(p+q+r)}(x)\\ &\quad-Z^{(p+q+r)}(x,\theta+\Phi(r))+\frac{p+q+r-\psi(\theta+\Phi(r))}{\Phi(p+q+r)-\theta-\Phi(r)}W^{(p+q+r)}(x)\\ &= Z^{(p+q+r)}(x,\Phi(p+r))\omega_1^-(0)+Z^{(p+q+r)}(x,\theta)-Z^{(p+q+r)}(x,\theta+\Phi(r)).\end{aligned}$$

Similarly,

$$\begin{aligned}\omega_2^-(x) &= \int_{-\infty}^0\mathbb{E}_x[e^{-(p+q)\tau_0^-};\tau_0^-<e_r,X_{\tau_0^-}\in dy]\mathbb{E}_y[e^{-p\tau_0^+};\tau_0^+<e_r]\omega_2^-(0)+\mathbb{E}_x[e^{-(p+q)e_r-\theta X(e_r)};e_r<\tau_0^-]\\ &= \frac{r}{q}\times\frac{\Phi(p+q+r)-\Phi(p+r)}{\theta+\Phi(p+q+r)}\left\{Z^{(p+q+r)}(x,\Phi(p+r))-\frac{q}{\Phi(p+q+r)-\Phi(p+r)}W^{(p+q+r)}(x)\right\}\\ &\quad+\frac{r}{\theta+\Phi(p+q+r)}W^{(p+q+r)}(x)-r\int_0^xe^{-\theta y}W^{(p+q+r)}(x-y)dy\\ &= \frac{r}{q}\times\frac{\Phi(p+q+r)-\Phi(p+r)}{\theta+\Phi(p+q+r)}Z^{(p+q+r)}(x,\Phi(p+r))-r\int_0^xe^{-\theta y}W^{(p+q+r)}(x-y)dy,\end{aligned}$$

where

$$\begin{aligned}&\mathbb{E}_x[e^{-(p+q)e_r-\theta X(e_r)};e_r<\tau_0^-]\\ &= r\int_0^\infty e^{-\theta y}\left(e^{-\Phi(p+q+r)y}W^{(p+q+r)}(x)-W^{(p+q+r)}(x-y)\right)dy\\ &= \frac{r}{\theta+\Phi(p+q+r)}W^{(p+q+r)}(x)-r\int_0^xe^{-\theta y}W^{(p+q+r)}(x-y)dy.\end{aligned}$$

We thus finish the proof. □

Using Theorem 3.2 we can find the probability that X hits 0 at time $T^-(r)$. It shows that $\mathbb{P}_x\{X_{T^-(r)} = 0\} > 0$ if and only if process X has a nontrivial Brownian component, which is similar to what happens at the first exit time.

Corollary 3.1. *For any $p, q, t > 0$ and $x \in \mathbb{R}$, we have*

$$\begin{aligned}&\mathbb{E}_x[e^{-pT^-(r)-q\int_0^{T^-(r)} 1_{X_s\geq 0}ds}; T^-(r) < e_r, X_{T^-(r)} = 0]\\ &= \frac{\sigma^2}{2}\Phi(r)Z^{(p+q+r)}(x, \Phi(p+r))\frac{1}{q}(\Phi(p+q+r) - \Phi(p+r)) - \frac{\sigma^2}{2}\Phi(r)W^{(p+q+r)}(x).\end{aligned} \tag{21}$$

In particular, for $\psi'(0+) < 0$,

$$\mathbb{E}_x[e^{-pT^-}; X_{T^-} = 0] = \frac{\sigma^2\Phi(0)}{2}\left(e^{\Phi(p)x}\Phi'(p) - W^{(p)}(x)\right).$$

Proof. Notice that

$$\begin{aligned}&\lim_{\theta\to\infty}\left(\frac{p+q+r-\psi(\theta+\Phi(r))}{\Phi(p+q+r)-\theta-\Phi(r)} - \frac{p+q+r-\psi(\theta)}{\Phi(p+q+r)-\theta}\right)\\ &= \lim_{\theta\to\infty}\frac{-\psi(\theta+\Phi(r))(\Phi(p+q+r)-\theta) + \psi(\theta)(\Phi(p+q+r)-\theta) + \Phi(r)(p+q+r-\psi(\theta))}{(\Phi(p+q+r)-\theta-\Phi(r))(\Phi(p+q+r)-\theta)}\\ &= -\frac{\sigma^2}{2}\Phi(p+q+r) + \sigma^2\Phi(r) + \frac{\sigma^2}{2}(\Phi(p+q+r) - \Phi(r))\\ &= \frac{\sigma^2}{2}\Phi(r),\end{aligned} \tag{22}$$

see Section 2.4 of [7].

In addition,

$$\begin{aligned}&\lim_{\theta\to\infty}\left(Z^{(p+q+r)}(x,\theta) - Z^{(p+q+r)}(x,\theta+\Phi(r))\right)\\ &= \lim_{\theta\to\infty}\left(\mathbb{E}_x e^{-(p+q+r)\tau_0^- + \theta X(\tau_0^-)}1_{\tau_0^-<\infty} + \frac{p+q+r-\psi(\theta)}{\Phi(p+q+r)-\theta}W^{(p+q+r)}(x)\right.\\ &\qquad\left. -\mathbb{E}_x e^{-(p+q+r)\tau_0^- + (\theta+\Phi(r))X(\tau_0^-)}1_{\tau_0^-<\infty} - \frac{p+q+r-\psi(\theta+\Phi(r))}{\Phi(p+q+r)-\theta-\Phi(r)}W^{(p+q+r)}(x)\right)\\ &= -\frac{\sigma^2}{2}\Phi(r)W^{(p+q+r)}(x).\end{aligned} \tag{23}$$

Then (21) follows from (18). □

Corollary 3.2. *For $\psi'(0+) < 0$ and $x \in \mathbb{R}$,*

$$\begin{aligned}&\mathbb{E}_x e^{-pT^- + \theta X_{T^-} - q\int_0^{T^-} 1_{X_s\geq 0}ds}\\ &= Z^{(p+q)}(x,\Phi(p))\frac{1}{q}(\Phi(p+q)-\Phi(p))\left(\frac{p+q-\psi(\theta+\Phi(0))}{\Phi(p+q)-\theta-\Phi(0)} - \frac{p+q-\psi(\theta)}{\Phi(p+q)-\theta}\right)\\ &\quad + Z^{(p+q)}(x,\theta) - Z^{(p+q)}(x,\theta+\Phi(0)).\end{aligned} \tag{24}$$

Remark 3.3. *For $x < 0$,*

$$\mathbb{E}_x e^{-pT^-(r)} = 1 - e^{\Phi(r)x} + e^{\Phi(p+r)x}\frac{\Phi'(p+r)}{\Phi(p+r)} \times \frac{p\Phi(r)}{\Phi(p+r)-\Phi(r)}.$$

For $x > 0$,

$$\begin{aligned}\mathbb{E}_x e^{-pT^-(r)} &= Z^{(p+r)}(x,\Phi(p+r))\frac{\Phi'(p+r)}{\Phi(p+r)} \times \frac{(p+r)\Phi(r) - r\Phi(p+r)}{\Phi(p+r)-\Phi(r)} + Z^{(p+r)}(x) - Z^{(p+r)}(x,\Phi(r)) \\ &\quad + \frac{r\Phi'(p+r)}{\Phi(p+r)}Z^{(p+r)}(x,\Phi(p+r)) - r\int_0^x W^{(p+r)}(x-y)dy \\ &= e^{\Phi(p+r)x}\frac{\Phi'(p+r)}{\Phi(p+r)} \times \frac{p\Phi(r)}{\Phi(p+r)-\Phi(r)} + \frac{r}{p+r} + \frac{p}{p+r}Z^{(p+r)}(x) - Z^{(p+r)}(x,\Phi(r)).\end{aligned}$$

We thus recover Theorem 2 of [2].

Remark 3.4. *For $\psi'(0+) < 0$ and $x \in \mathbb{R}$, letting $q, \theta, r \to 0+$ we have*

$$\mathbb{E}_x e^{-pT^-} = Z^{(p)}(x,\Phi(p))\frac{p\Phi(0)\Phi'(p)}{\Phi(p)(\Phi(p)-\Phi(0))} + Z^{(p)}(x) - Z^{(p)}(x,\Phi(0));$$

letting $p, q, r \to 0+$, we have

$$\mathbb{E}_x e^{\theta X_{T^-}} = e^{\Phi(0)x}\Phi'(0)\left(\frac{\psi(\theta+\Phi(0))}{\theta} + \frac{\psi(\theta)}{\Phi(0)-\theta}\right) + Z(x,\theta) - Z(x,\theta+\Phi(0));$$

letting $p, \theta, r \to 0+$, we have

$$\begin{aligned}\mathbb{E}_x e^{-q\int_0^\infty 1_{X_s>0}ds} &= Z^{(q)}(x,\Phi(0))\frac{\Phi(0)}{\Phi(q)} + Z^{(q)}(x,0) - Z^{(q)}(x,\Phi(0)) \\ &= Z^{(q)}(x,\Phi(0))\left(\frac{\Phi(0)}{\Phi(q)} - 1\right) + Z^{(q)}(x).\end{aligned}$$

which recovers Corollary 3 (iv) of [9].

We now consider some examples for more explicit results. Let $X_t = -\mu t + B_t$ with $\mu > 0$ and Brownain motion B. Then $T^- < \infty$ a.s.,

$$\psi(\lambda) = -\mu\lambda + \frac{\lambda^2}{2},\ \Phi(q) = \sqrt{\mu^2+2q} + \mu,$$

$$W^{(q)}(x) = \frac{1}{\sqrt{\mu^2+2q}}\left(e^{(\sqrt{\mu^2+2q}+\mu)x} - e^{-(\sqrt{\mu^2+2q}-\mu)x}\right)1_{x\geq 0}$$

and

$$W(x) = \frac{1}{\mu}(e^{2\mu x} - 1)1_{x\geq 0}.$$

By Corollary 3.1,

$$
\begin{aligned}
&\mathbb{E}_x[e^{-pT^-}; X_{T^-}=0] \\
&= \frac{\Phi(0)}{2}\left(e^{\Phi(p)x}\Phi'(p) - W^{(p)}(x)\right) \\
&= \mu\left(\frac{1}{\sqrt{\mu^2+2p}}e^{(\sqrt{\mu^2+2p}+\mu)x} - \frac{1}{\sqrt{\mu^2+2p}}\left(e^{(\sqrt{\mu^2+2p}+\mu)x} - e^{-(\sqrt{\mu^2+2p}-\mu)x}\right)1_{x\geq 0}\right) \\
&= \frac{\mu}{\sqrt{\mu^2+2p}}\left(e^{(\sqrt{\mu^2+2p}+\mu)x}1_{x<0} + e^{-(\sqrt{\mu^2+2p}-\mu)x}1_{x\geq 0}\right).
\end{aligned} \tag{25}
$$

It follows that

$$
\mathbb{P}_x\{X_{T^-}=0\} = e^{2\mu x}1_{x<0} + 1_{x\geq 0} = \mathbb{P}_x\{\tau_0^+ < \infty\}1_{x<0} + 1_{x\geq 0},
$$

which is expected intuitively.

Let X be a spectrally negative α-stable process with a negative drift whose Laplace exponent is given by $\psi(\lambda) = -\mu\lambda + \lambda^\alpha$ for $\mu > 0$ and $1 < \alpha < 2$. Then $\Phi(0) = \mu^{\frac{1}{\alpha-1}}$ and by Remark 3.4.

$$
\begin{aligned}
\mathbb{E}e^{\theta X_{T^-}} &= \frac{1}{\psi'(\Phi(0))}\left(\frac{\psi(\theta+\Phi(0))}{\theta} + \frac{\psi(\theta)}{\Phi(0)-\theta}\right) \\
&= \frac{1}{(\alpha-1)\mu}\left(\frac{-\mu\theta - \mu^{\frac{\alpha}{\alpha-1}} + (\theta+\mu^{\frac{1}{\alpha-1}})^\alpha}{\theta} + \frac{-\mu\theta+\theta^\alpha}{\mu^{\frac{1}{\alpha-1}}-\theta}\right).
\end{aligned} \tag{26}
$$

It is not hard to see that

$$
\lim_{\theta\to\infty}\mathbb{E}e^{\theta X_{T^-}} = 0,
$$

i.e. $\mathbb{P}\{X_{T^-}=0\} = 0$ which agrees with Corollary 3.1.

Let X be a compound Poisson process with a positive drift μ and negative exponential jumps with common rate ρ arriving according to a Poisson process with rate a. We further assume that $\mu - \frac{a}{\rho} < 0$ so that the last exit time is finite. Then

$$
\psi(\lambda) = \mu\lambda - \frac{a\lambda}{\rho+\lambda}, \quad \Phi(0) = \frac{a}{\mu} - \rho,
$$

$$
\psi'(\Phi(0)) = \mu - \frac{a\rho}{\left(\rho + \frac{a}{\mu} - \rho\right)^2} = \mu - \frac{\rho\mu^2}{a}
$$

and

$$
\begin{aligned}
\psi(\theta+\Phi(0)) &= \mu(\theta + \frac{a}{\mu} - \rho) - \frac{a(\theta+\frac{a}{\mu}-\rho)}{\rho+\theta+\frac{a}{\mu}-\rho} \\
&= (\theta + \frac{a}{\mu} - \rho)\frac{\mu\theta}{\theta+\frac{a}{\mu}}.
\end{aligned} \tag{27}
$$

It follows that

$$\begin{aligned}\mathbb{E}e^{\theta X_{T^-}} &= \frac{1}{\psi'(\Phi(0))}\left(\frac{\psi(\theta+\Phi(0))}{\theta}+\frac{\psi(\theta)}{\Phi(0)-\theta}\right)\\ &= \frac{1}{\mu-\frac{\rho\mu^2}{a}}\left((\theta+\frac{a}{\mu}-\rho)\frac{\mu}{\theta+\frac{a}{\mu}}+\frac{\mu\theta-\frac{a\theta}{\rho+\theta}}{\frac{a}{\mu}-\rho-\theta}\right)\\ &= \frac{a\rho}{(\rho+\theta)(\mu\theta+a)}\\ &= \frac{a}{a-\mu\rho}\times\frac{\rho}{\rho+\theta}-\frac{\mu\rho}{a-\mu\rho}\times\frac{a}{a+\mu\theta}.\end{aligned} \tag{28}$$

Therefore, X_{T^-} has a density

$$\left(\frac{a}{a-\mu\rho}\rho e^{\rho x}-\frac{\mu\rho}{a-\mu\rho}\times\frac{a}{\mu}e^{\frac{ax}{\mu}}\right)1_{x<0}=\frac{a\rho}{a-\mu\rho}\left(e^{\rho x}-e^{\frac{ax}{\mu}}\right)1_{x<0}.$$

It is remarkable that X_{T^-} does not follow an exponential distribution. In addition, compared with the exponential random variable e_ρ, $-X_{T^-}$ has a better chance to take large values. Notice that X_{T^-} does not have the same density of e_ρ, the Laplace transform of X_{T^-} can not be directly obtained from the distribution of X_{T^--} due to the the fact that T^- is not a stopping time.

References

[1] H. Albrecher, J. Ivanovs and X. Zhou. Exit identities for a Lévy processes observed at Poisson arrival times. *Bernoulli 22 (2016), 1364-1382.*

[2] E. J. Baurdoux. Last exit before an exponential time for spctrally negativw Lévy processes. *Journal of Applied Probability 46 (2009), 542-588*

[3] S. N. Chiu and C. Yin. Passage times for a spectrally negative Lévy process with applications to risk theory. *Bernoulli 11 (2005), 511-522.*

[4] H.U. Gerber. When does the surplus reach a given target? *Insurance, Mathematics & Econoimics 9 (1990), 115-119.*

[5] B. Li and C. Cai. Occupation times of intervals until last passage times for spectrally negative Lévy processes. Preprint. arXiv 1605.07709.

[6] Y. Li and X. Zhou. On pre-exit joint occupation times for spectrally negative Lévy processes. *Statistics and Probability Letters 94 (2014), 48-55.*

[7] A. Kuznetsov, A.E. Kyprianou, V. Rivero. The theory of scale functions for spectrally negative Lévy processes. In: Lévy Matters, in: Springer Lecture Notes in Mathematics, 2013, pp. 97-186.

[8] A. E. Kyprianou. Introductory Lectures on Fluctuations of Lévy processes with Applications. 2nd edition. Universitext. Springer-Verlag, Berlin, 2014.

[9] R. Loeffen, J.-F. Renaud and X. Zhou. Occupation times of intervals until first passage times for spectrally negative Lévy processes. *Stochastic Processes and their Applications 124 (2014) 1408-1435.*

[10] X. Zhou. Exit problems for spectrally negative Lévy processes reflected at either the supremum or the infimum. *Journal of Applied Probability 44 (2007), 1012–1030.*